

# A dynamic model of the wormhole and the Multiverse model

A A Shatskii, I D Novikov, N S Kardashev

DOI: 10.1070/PU2008v051n05ABEH006581

## Contents

1. Introduction	457
2. Equations of motion	458
3. Integration	458
4. Initial conditions	459
5. Solution without the $\Lambda$ term	460
6. Solution $s = 0$ with the $\Lambda$ term	461
7. Static solution with the $\Lambda$ term; a model of the Multiverse	461
8. Conclusion	462
9. Appendices	462
References	463

**Abstract.** An analytic solution methodology for general relativity (GR) equations describing the hypothetical phenomenon of wormholes is presented and the analysis of wormholes in terms of their physical properties is discussed. An analytic solution of the GR equations for static and dynamic spherically symmetric wormholes is given. The dynamic solution generally describes a ‘traversable’ wormhole, i.e., one allowing matter, energy, and information to pass through it. It is shown how the energy–momentum tensor of matter in a wormhole can be represented in a form allowing the GR equations to be solved analytically, which has a crucial methodological importance for analyzing the properties of the solution obtained. The energy–momentum tensor of wormhole matter is represented as a superposition of a spherically symmetric magnetic (or electric) field and negative-density dust matter, serving as exotic matter necessary for a ‘traversable’ wormhole to exist. The dynamics of the model are investigated. A similar model is considered (and analyzed in terms of inflation) for the Einstein equations with a  $\Lambda$  term. Superposing enough dust matter, a magnetic field, and a  $\Lambda$  term can produce a static solution, which turns out to be a spherical Multiverse model with an infinite number of wormhole-connected spherical universes. This Multiverse can have its total energy positive everywhere in space, and in addition can be out of equilibrium (i.e., dynamic).

## 1. Introduction

In the last few years, the problem of wormholes (WHs) in General Relativity (GR) has been discussed in numerous papers (see, e.g., [1–19]). This problem is relevant to both theoretical physics and astrophysics.

A wormhole is a hypothetical object described by a non-singular solution of the Einstein equations with two large (or infinite) space–time regions connected by a throat. The two large space–time regions can be located in one universe or belong to different universes in the Multiverse model (see [18]). In the last case, ‘traversable’ WHs afford a unique opportunity to explore other universes.

In the present paper, we analytically study the dynamics of a spherical model of a nonequilibrium WH filled with matter. This matter consists of a magnetic (or electric) field and dust with negative mass density. The obtained solution therefore generalizes the Tolman solution [20] for a model with a centrally symmetric electromagnetic field. As we see below, this generalization leads to essentially new and important solutions. We use the method of calculations of physical quantities in the frame comoving with matter. This method was elaborated by Zel’manov [21] and applied by Novikov [22, 23] (see also [16]) to spherically symmetric problems and was later utilized and developed for many similar problems.

As the initial model for a WH, we use a static model in which gravitational accelerations are everywhere identically zero. Hence, the effective masses of both WH mouths vanish, although the geometry of three-dimensional space is certainly non-Euclidean. Such a model is considered in [17, 19], where all matter is represented by a gravitating scalar field. We change the scalar field into a superposition of an electromagnetic field and dust matter with negative energy density, which turns out to be a methodologically important development and generalization of these models. This allows us to apply methods of Tolman’s problem to calculate the model (see [20, 24, 25]); these methods were generalized and further developed by Shatskii (see [26, 27]).

The results and methods in this paper can be used in discussing the possible evolution of a WH in different models

A A Shatskii, N S Kardashev Astro-Space Center, Lebedev Physical Institute, Russian Academy of Sciences,

Profsoyuznaya ul. 84/32, 117997 Moscow, Russian Federation  
Tel. (7-495) 333 33 66; 333 21 89; 334 85 06. Fax (7-495) 333 23 78  
E-mail: shatskiy@asc.rssi.ru; nkardash@asc.rssi.ru

I D Novikov Astro-Space Center, Lebedev Physical Institute, Russian Academy of Sciences,

Profsoyuznaya ul. 84/32, 117997 Moscow, Russian Federation;  
Niels Bohr Institute, Blegdamsvej 17, DK-2100 Copenhagen, Denmark;  
Russian Research Center Kurchatov Institute,  
pl. Kurchatova 1, 123182 Moscow, Russian Federation

Received 12 February 2008, revised 31 March 2008

*Uspekhi Fizicheskikh Nauk* 178 (5) 481–488 (2008)

DOI: 10.3367/UFNr.0178.200805c.0481

Translated by K A Postnov; edited by A M Semikhatov

of the Universe. To further generalize and develop this method, we introduce the cosmological  $\Lambda$  term into the model. This allows obtaining a principally new solution (see Section 7) for a static spherical model of the Multiverse. This model includes a infinite number of spherical worlds connected by throats. To our knowledge, this is the first analytic model of this type.

In the obtained solution, the Multiverse can have its total energy density positive everywhere in space. In addition, this solution can be generalized by the same method to the case of a dynamic model shifted from equilibrium by an excess (or shortage) of dust or by the  $\Lambda$  term.

To conclude this section, we make a methodological remark. Nonstationary problems in GR are principally different from those in flat space–time. In particular, the effects of the motion (dynamics) of matter in GR are accompanied by the dynamic effects of reference frame deformations (in our method, the comoving dust frame). Usually, it is very difficult to distinguish these effects. In this paper, the term ‘motion’ is used for a superposition of the dynamics and deformations.

## 2. Equations of motion

The simplest spherically symmetric solution of the Einstein equations for a static WH has the form (see [17])<sup>1</sup>

$$ds^2 = dt^2 - dR^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2),$$

$$r^2(R) = q^2 + R^2. \tag{1}$$

It follows that  $4\pi r^2$  represents the area of a sphere around the center of the system. The minimum allowed radius of this WH is  $r_0 = q$  (the radius of the throat).<sup>2</sup>

Usually, this solution is related to the energy–momentum tensor of a scalar field (see [17, 19]). In the present paper, we suggest another interpretation of this solution related to a different representation of the energy–momentum tensor. This new interpretation of solution (1) can correspond to the energy–momentum tensor represented as

$$T_m^n = \begin{pmatrix} -\frac{q^2}{8\pi r^4} & 0 & 0 & 0 \\ 0 & +\frac{q^2}{8\pi r^4} & 0 & 0 \\ 0 & 0 & -\frac{q^2}{8\pi r^4} & 0 \\ 0 & 0 & 0 & -\frac{q^2}{8\pi r^4} \end{pmatrix}$$

$$= \begin{pmatrix} +\frac{q^2}{8\pi r^4} & 0 & 0 & 0 \\ 0 & +\frac{q^2}{8\pi r^4} & 0 & 0 \\ 0 & 0 & -\frac{q^2}{8\pi r^4} & 0 \\ 0 & 0 & 0 & -\frac{q^2}{8\pi r^4} \end{pmatrix}$$

$$+ \begin{pmatrix} -\frac{q^2}{4\pi r^4} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \tag{2}$$

<sup>1</sup> We use units where  $c = 1$  and  $G = 1$ .

<sup>2</sup> Here and below, all values at the throat have a subscript 0.

The first term in the right-hand side corresponds to the energy–momentum tensor of a static electric (or magnetic) field with the charge  $q$  and the second term describes dust matter with the negative energy density

$$\varepsilon_d = -\frac{q^2}{4\pi r^4}. \tag{3}$$

This possible representation of the tensor  $T_m^n$  is different from its equivalent representation as the energy–momentum tensor of a scalar field adopted in previous papers [17–30].

An excess (or shortage) of dust (relative to  $\varepsilon_d$ ) is not in equilibrium and initiates motion. Because dust layers are relatively independent, it is possible to integrate the equations of motion for dust in a way similar to the solution of Tolman’s problem. In essence, it is the same as Tolman’s problem in a centrally symmetric and static electric (or magnetic) field for uncharged dust.

We seek the metric tensor of the solution in the form<sup>3</sup>

$$ds^2 = dt^2 - \exp(\lambda) dR^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2), \tag{4}$$

where  $r^2$  and  $\exp(\lambda)$  are functions of both  $R$  and  $t$ . Below, we justify this choice of the metric.

We consider the problem in the presence of a  $\Lambda$  term. The Einstein equations corresponding to metric (4) can be written as<sup>4</sup>

$$8\pi T_t^t = 8\pi\varepsilon + \frac{q^2}{r^4} + \Lambda$$

$$= -\exp(-\lambda) \left( \frac{2rr_{,RR} + r_{,R}^2 - rr_{,R}\lambda_{,R}}{r^2} + \frac{rr_{,t}\lambda_{,t} + r_{,t}^2}{r^2} + \frac{1}{r^2} \right), \tag{5}$$

$$8\pi T_R^R = \frac{q^2}{r^4} + \Lambda = \frac{2rr_{,tt} + r_{,t}^2}{r^2} - \frac{\exp(-\lambda)r_{,R}^2}{r^2} + \frac{1}{r^2}, \tag{6}$$

$$8\pi T_t^R = 0 = \exp(-\lambda) \frac{2r_{,Rt} - r_{,R}\lambda_{,t}}{r}, \tag{7}$$

$$8\pi T_\theta^\theta = 8\pi T_\varphi^\varphi = -\frac{q^2}{r^4} + \Lambda$$

$$= -\frac{\exp(-\lambda)(2r_{,RR} - r_{,R}\lambda_{,R}) - 2r_{,tt} - r\lambda_{,tt} - r\lambda_{,t}^2/2 - r_{,t}\lambda_{,t}}{2r}. \tag{8}$$

Thus, the energy–momentum tensor includes three types of matter:

- a centrally symmetric electric (or magnetic) field,
- a cosmological  $\Lambda$  term, and
- dust matter with the density  $\varepsilon$ .<sup>5</sup>

## 3. Integration

Integrating Eqn (7) with respect to time, we obtain

$$\exp(-\lambda) r_{,R}^2 = F_1(R). \tag{9}$$

<sup>3</sup> Because dust is pressureless, we can choose the synchronous comoving frame (with the time metric component equal to 1).

<sup>4</sup> See, e.g., [31] for the derivation (problem 5 in § 100).

<sup>5</sup> In the absence of the  $\Lambda$  term, solution (4) coincides with (1) only for the density  $\varepsilon$  equal to  $\varepsilon_d$  [see (3)].

At the throat, the condition  $r_{,R} = 0$  must be satisfied, and therefore

$$F_1(0) = 0. \tag{10}$$

By substituting (9) in Eqn (6), we obtain

$$\frac{q^2}{r^2} + Ar^2 = \frac{(rr_{,t})_{,t}}{r_{,t}} - F_1 + 1. \tag{11}$$

Integrating this equation with respect to time yields

$$\frac{q^2}{r} - \frac{Ar^3}{3} + rr_{,t}^2 + (1 - F_1)r = F_2(R). \tag{12}$$

The functions  $F_1(R)$  and  $F_2(R)$  determine the initial conditions for the velocity and dust density distributions.

With  $\lambda_{,t}$  and  $\lambda_{,tt}$  expressed from Eqn (7), Eqn (8) can be rewritten as

$$\left[ \frac{q^2}{r^2} + Ar^2 + \exp(-\lambda) r_{,R}^2 - r_{,t}^2 - 2rr_{,tt} \right]_{,R} = 0. \tag{13}$$

When Eqn (6) is satisfied, integrating Eqn (13) yields an identity. *This identity is the consequence of setting the time component of metric (4) to unity, which justifies this choice of the metric.*

It is worth noting from the methodological standpoint that this form of the metric for dust is obvious because dust is pressureless and hence the comoving frame is simultaneously a synchronous one. But this is not so obvious when a spherically symmetric electromagnetic field is added, and the obtained identity is a consequence of the absence of interaction between uncharged dust and the electromagnetic field. Moreover, the Lorentz transformation along the magnetic field lines does not change this field, which justifies the use of the dust-comoving reference frame. For motion in this frame (along the field), the electromagnetic field remains invariant. If the metric could not be represented in form (4), further calculations would be impossible. It is this nonobvious fact that allowed us to successfully develop our method and to obtain all important results.

Multiplying Eqn (5) by  $r^2 r_{,R}$  and expressing  $\lambda_{,t}$  from Eqn (7), we obtain

$$8\pi\epsilon r^2 r_{,R} + \left[ \frac{Ar^3}{3} - \frac{q^2}{r} - r(1 - F_1) - rr_{,t}^2 \right]_{,R} = 0. \tag{14}$$

Integrating this with respect to  $R$  from 0 to  $R$  with account for (10) yields

$$\int_0^R 8\pi\epsilon r^2 r_{,R} dR - \frac{q^2}{r} + \frac{Ar^3}{3} - r(1 - F_1) - rr_{,t}^2 = -\frac{q^2}{r_0} + \frac{Ar_0^3}{3} - r_0 - r_0 r_{0,t}^2. \tag{15}$$

Here, we took into account that the function  $r(R = 0) = r_0(t)$  can be time-dependent.

Expressing  $rr_{,t}^2$  from (12) and substituting it in (15), we obtain

$$\int_0^R 8\pi\epsilon r^2 r_{,R} dR = F_2(R) - \frac{q^2}{r_0} - r_0 + \frac{Ar_0^3}{3} - r_0 r_{0,t}^2. \tag{16}$$

The derivative of this expression with respect to  $R$  yields one more integral of motion (which expresses mass conservation in the comoving volume):

$$8\pi\epsilon r^2 r_{,R} = \frac{dF_2}{dR}. \tag{17}$$

The Bianchi identities<sup>6</sup>

$$T_{m;n}^n = \frac{1}{\sqrt{-g}} \frac{\partial(T_m^n \sqrt{-g})}{\partial x^n} - \frac{T^{nl}}{2} \frac{\partial g_{nl}}{\partial x^m} = 0 \tag{18}$$

are contained in the Einstein equations, their components  $m = t, \theta, \varphi$  vanish identically, and the component  $m = R$  yields the result in (17), which is an analogue of the integral for the pure-dust solution (see [31], § 103) in the theory of the evolution of a dust cloud. The explicit form of the function  $\exp(\lambda)(R, t) = r_{,R}^2/F_1$  and an implicit form of the function  $r^2(R, t)$  are obtained in Section 5.

### 4. Initial conditions

Below, we use the index ‘i’ to denote all quantities at the initial instant  $t = 0$ . The velocity distribution is set to zero initially:

$$r_{,t}^2|_{t=0} = 0. \tag{19}$$

A nonzero initial velocity is considered in Appendix I. We note that in all cases, the choice of initial conditions is not arbitrary and must satisfy constraints imposed by Eqns (5) and (7). These equations do not contain second time derivatives and, as is well known (see [29]), under the condition that  $t = \text{const}$  at the initial cross section, they are automatically satisfied for all values by virtue of the other Einstein equations.

We choose the initial coordinate scale in (4) such that

$$r_i = \sqrt{q^2 + R^2}, \quad \epsilon_{di} \equiv \epsilon_d(r_i) = -\frac{q^2}{4\pi r_i^4}, \tag{20}$$

where  $r_i(R)$  coincides with (1) and Eqn (10) is automatically satisfied. We set

$$s(r_i) \equiv \frac{1}{r_i} \int_0^R 8\pi(\epsilon_i - \epsilon_{di}) r_i^2 r_{i,R} dR = \frac{1}{r_i} \int_q^{r_i} 8\pi\epsilon_i r_i^2 dr_i + 2\frac{q}{r_i} \left(1 - \frac{q}{r_i}\right). \tag{21}$$

The quantity  $s$  has the meaning of the doubled Newtonian potential of dust matter excessive relative to  $\epsilon_d$  [see (20)]. The relation of this potential to the functions  $F_1$  and  $F_2$  follows from (15) and (16) at the initial instant:

$$F_1 = 1 - s - \frac{q^2}{r_i^2} - \frac{A}{3} \left( r_i^2 - \frac{q^3}{r_i} \right), \tag{22}$$

$$F_2 = r_i \left( s + 2\frac{q^2}{r_i^2} - \frac{Aq^3}{3r_i} \right). \tag{23}$$

Equation (9) implies that  $F_1 \geq 0$ . According to (22), this imposes constraints on the distribution  $s(R)$ .

<sup>6</sup> Here,  $g$  is the determinant of the metric tensor.

From Eqn (6), we can find  $r_{,tt}$  at the initial instant:

$$r_{,tt} = -\frac{s}{2r_i} + \frac{Ar_i}{3} + \frac{Aq^3}{6r_i^2} \quad \text{for } t = 0. \quad (24)$$

Equation (24) shows that for  $A = 0$ , the dynamics are absent in the region where  $s = 0$  (at least up to the instant of intersection of the dust layers, when the model considered becomes inapplicable).

## 5. Solution without the $A$ term

We consider the case  $A = 0$  in more detail.

For  $A = 0$ , it follows from Eqn (24) that the WH throat size remains constant [ $r_0(t) = q$ ] in the whole region of the allowed values of  $R$  and  $t$  for the sought solution, until matter starts flowing through the throat. In addition, expression (24) implies that the dynamics of matter in a WH depend only on the internal layers of matter (with smaller values of  $r$ ) and are independent of the external layers. Therefore, matter on one side of the throat does not influence matter on the opposite side (unless there is matter flowing through the throat or layer intersection occurs).

It can be verified directly from (24) that at the initial instant, the acceleration  $r_{,tt}$  is negative for  $s > 0$ . Hence, the excessive mass of dust is to collapse.

We introduce the definition of the visibility horizon.

*The criterion of the absence of a horizon (visibility) during the collapse is given by*

$$V < 1 \quad (25)$$

for the dust velocity [see also [23, 26, 27, 32] and (12)], where the dust velocity  $V$  is determined as

$$V^2 \equiv \frac{r_{,t}^2 \exp(\lambda)}{r_{,R}^2} = \frac{r_{,t}^2}{F_1} = \frac{(s + 2q^2/r_i^2)r_i/r - s - q^2/r_i^2 - q^2/r^2}{1 - s - q^2/r_i^2}. \quad (26)$$

For  $q = 0$ , this result coincides with the solution for collapsing dust (see [26]).

To avoid a 0/0 ambiguity for the function  $V^2$  at the throat at the initial instant ( $F_1[r_i \rightarrow r_0] \rightarrow 0$ ), it is necessary to ensure a more rapid convergence to zero of the function  $r_{,t}^2(t = 0, r_i \rightarrow r_0)$  than the function  $F_1(r_i \rightarrow r_0)$  (see Appendix I).

The law of motion of a dust layer can be found from (12):

$$t = \int_{r_i}^r \frac{-r dr}{\sqrt{(F_1 - 1)r^2 + F_2r - q^2}} = \int_{r_i}^r \frac{-r dr}{r_i \sqrt{(1 - r/r_i)[(s + q^2/r_i^2)r/r_i - q^2/r_i^2]}}. \quad (27)$$

This quadrature can be elementarily integrated:

$$t = r_i \frac{\sqrt{(1 - r/r_i)[(s + q^2/r_i^2)r/r_i - q^2/r_i^2]}}{s + q^2/r_i^2} + r_i \frac{s + 2q^2/r_i^2}{2(s + q^2/r_i^2)^{3/2}} \arccos \left[ 1 - \frac{2(s + q^2/r_i^2)(1 - r/r_i)}{s} \right], \quad (28)$$

or in another form,

$$t = r_i \left[ \frac{\sqrt{(1 - r/r_i)[(s + q^2/r_i^2)r/r_i - q^2/r_i^2]}}{s + q^2/r_i^2} + \frac{s + 2q^2/r_i^2}{(s + q^2/r_i^2)^{3/2}} \arctan \sqrt{\frac{(s + q^2/r_i^2)(1 - r/r_i)}{(s + q^2/r_i^2)r/r_i - q^2/r_i^2}} \right]. \quad (29)$$

*Hence, the motion of dust layers depends on the excessive potential  $s$  and the initial distance from the WH throat:  $r_i/q$ .*

However, this model is physically correct only until a possible intersection of dust layers. This is because after the intersection, an incoming energy flux appears in the matter-moving frame (from the intersected matter layers), which has not been taken into account in the energy–momentum tensor. The intersection of adjacent matter layers corresponds to an infinite energy density. This corresponds to the condition  $r_{,R} = 0$  (or  $dr/dr_i = 0$ ) [see (20)]. Differentiation of (29) with respect to  $r_i$  allows explicitly determining the function  $dr/dr_i(r, s, q)$ . Differentiation of the left-hand side of (29) yields zero. We let a prime denote the derivative with respect to  $r_i$ , express all distances in the units of units  $r_i$ , and omit intermediate bulky calculations; the final expression for this function is given by

$$r' = 1 + \frac{(1 - r)\{s(s' - 2q^2)[2q^2 - r(s + q^2)] - q^2(s + 2q^2)(2s' + s)\}}{2rs(s + q^2)^2} + \frac{[2s(s + q^2) + (2q^2 - s')(s + 4q^2)]\sqrt{(1 - r)[(s + q^2)r - q^2]}}{2r(s + q^2)^{5/2}} \times \arctan \sqrt{\frac{(s + q^2)(1 - r)}{(s + q^2)r - q^2}}. \quad (30)$$

It is impossible to use this formula to determine the instant of intersection of nonadjacent layers because the intersection with nonadjacent layers does not lead to any irregularities for dust. Clearly, the intersection of nonadjacent layers occurs after the intersection of adjacent ones and cannot propagate with a superluminal velocity. Therefore, after constructing diagrams  $t(r_i)$  for light cones and the curve corresponding to the intersection of adjacent layers ( $r' = 0$ ), we can determine regions that are definitely free from layer intersection (see the Figure).

Using formula (26), we can reach several important conclusions.

1. The formation of a horizon is possible only for a sufficiently large parameter  $s$ :

$$s > 2 \frac{q}{r_i} \left( 1 - \frac{q}{r_i} \right). \quad (31)$$

After the appearance of a horizon, the WH becomes ‘nontraversable.’ However, dust layers can already start intersecting before the horizon is reached.

2. Values of the function  $t(r, r_i)$  are bounded by the maximum time  $t_{\text{stop}}$  (from the beginning of motion until the stop). In (28), this time corresponds to the argument of arccos being minus unity (or to the zero value of the square root expression in the same formula):

$$t_{\text{stop}} = \frac{\pi r_i (s + 2q^2/r_i^2)}{2(s + q^2/r_i^2)^{3/2}}. \quad (32)$$

The time  $t_{\text{stop}}$  corresponds to the deviation  $\Delta r \approx sr_i^3/q^2$  from the initial position. For small deviations (corresponding to small values of  $s$ ), harmonic oscillations with the period  $T = 2t_{\text{stop}} \approx 2\pi r_i^2/q$  occur (see Appendix II).

3. The existence of the second (smaller) root of the equation  $V^2(r) = 1$  implies the appearance of the second (inner) horizon in the system (see the Figure).

### 6. Solution $s = 0$ with the $\Lambda$ term

We consider the case where  $s = 0$ ; Eqn (6) then becomes

$$2rr_{,tt} = \Lambda r^2 + q^2 \left( \frac{1}{r^2} - \frac{1}{r_i^2} \right) - \frac{\Lambda}{3} \left( r_i^2 - \frac{q^3}{r_i} \right) - r_{,t}^2, \quad (33)$$

and Eqn (12) takes the form


$$r_{,t}^2 = (r - r_i) \left[ -\frac{q^2}{r^2 r_i^2} (r - r_i) + \frac{\Lambda}{3} \left( r + r_i + \frac{q^3}{rr_i} \right) \right]. \quad (34)$$

These two equations imply that at  $s = 0$ , matter starts expanding from the rest state (inflation due to the  $\Lambda$  term). If the dimensionless parameter

$$a \equiv \Lambda q^2 \quad (35)$$

exceeds some critical value  $a_{\text{cr}}$ , the inflation continues unlimitedly in the entire volume ( $q \leq r_i \leq r < \infty$ ) until an outer horizon forms,  $V^2(r) = 1$ . The value  $a_{\text{cr}}$  is found from the condition of the maximum of the expression for the parameter  $a$ , which is obtained by equating the second factor (in square brackets) in Eqn (34) to zero:

$$a_{\text{cr}} = \max \left[ \frac{3\tilde{q}^4(1 - \tilde{r})\tilde{r}^2}{1 + \tilde{r} + \tilde{r}^2\tilde{q}^3} \right] \approx 0.22, \quad \tilde{r} \equiv \frac{r_i}{r}, \quad \tilde{q} \equiv \frac{q}{r_i}. \quad (36)$$



**Figure.** Diagrams  $t(r_i)$ : 1, for light geodesics; 2, for  $r = \text{const}$ ; 3, for the visibility horizon and the inner horizon; 4, for the limiting stopping time; 5, for the throat  $r = q$ ; 6, for the intersection times of adjacent dust layers; 7, the dependence  $s(r_i) = 0.7(r_i/q - 1) \exp(2 - r_i/q)$ ; the scale for  $s$  (vertical axis) coincides with the time scale  $t$ .

For  $a < a_{\text{cr}}$ , a region appears where the inflation stops. This region starts emerging at the throat and extends toward larger values of  $r_i$  as the parameter  $a$  decreases.

As  $r/r_i \rightarrow \infty$ , the inflation occurs exponentially:  $r \propto \exp(t\sqrt{\Lambda/3})$ .

The function  $r'$  can be expressed in quadratures (see Appendix III):

$$r' = 1 + \sqrt{r - r_i} \exp \left( - \int_{r_i}^r P_2(x) dx \right) \int_{r_i}^r \frac{Q_2(r) - P_2(r)}{\sqrt{r - r_i}} \times \exp \left( \int_{r_i}^r P_2(x) dx \right) dr, \\ P_2 = \frac{r_i^2/(2r^2) - r_i^2(r - r_i)/r^3 - a/2[1 - q^3/(r^2 r_i)]}{-r_i^2(r - r_i)/r^2 + a[r + r_i + q^3/(rr_i)]}, \\ Q_2 - P_2 = \frac{a + (r + r_i)[r_i(r - r_i)/r^3 - aq^3/(2r^2 r_i^2)]}{-r_i^2(r - r_i)/r^2 + a[r + r_i + q^3/(rr_i)]}. \quad (37)$$

Using this equation, we can find regions where the inflation occurs without dust layer intersections.

### 7. Static solution with the $\Lambda$ term; a model of the Multiverse

Dust with positive mass excessive relative to  $\varepsilon_d$  initially accelerates toward the center (without the  $\Lambda$  term). The  $\Lambda$  term without excessive dust leads to the original inflationary solution [see (24)]. At the WH throat, the potential  $s = 0$ , and the  $\Lambda$  term at the throat provides nonzero matter acceleration. This contribution cannot be compensated at or near the throat [see (6)], where  $r_{,t} = r_{,tt} = 0$ , and  $r = q$ . Hence, there is no static solution for a WH with the throat radius  $r_0 = q$  and the  $\Lambda$  term.

Nevertheless, a static solution exists for a WH with the  $\Lambda$  term and  $r_0 \neq q$ . This solution can be easily derived from Eqns (5), (6), and (8) (with  $r_{,t} = 0$  and  $r_{,tt} = 0$ ).

The metric of a static WH is determined by expression (1), taking into account that  $r^2(R) \neq q^2 + R^2$ .

For (6), we obtain  $r_{,R}^2 = 1 - q^2/r^2 - \Lambda r^2$  and easily find the expression for the throat radius  $r_0$ ,

$$\Lambda r_0^4 - r_0^2 + q^2 = 0 \Rightarrow r_0^2 = \frac{1 - \sqrt{1 - 4\Lambda q^2}}{2\Lambda} \\ \rightarrow q^2(1 + \Lambda q^2) \text{ for } \Lambda q^2 \rightarrow 0, \quad (38)$$

and the dependence  $r(R)$ :

$$r^2(R) = \frac{1 - \sqrt{1 - 4\Lambda q^2} \cos(2\sqrt{\Lambda}R)}{2\Lambda}. \quad (39)$$

The distribution of  $\varepsilon$  for this solution is determined<sup>7</sup> from (5):

$$\varepsilon = \frac{\Lambda}{4\pi} - \frac{q^2}{4\pi r^4}. \quad (40)$$

Thus, for the static solution with the  $\Lambda$  term, the throat radius and dust density are larger than without the  $\Lambda$  term.

<sup>7</sup> Here, the unknown value  $r_{,RR} = q^2/r^3 - \Lambda r$  can be obtained from expression (8) or by directly differentiating Eqn (39).

The total energy density for the static solution with the  $\Lambda$  term is

$$T'_t = \frac{3\Lambda}{8\pi} - \frac{q^2}{8\pi r^4}. \tag{41}$$

The condition for the value  $T'_t$  to be nonnegative everywhere in space (for  $r \geq r_0$ ) has the form

$$T'_t \geq 0 \text{ for } a = \Lambda q^2 \geq \frac{3}{16}. \tag{42}$$

Therefore, in the presence of the  $\Lambda$  term, solutions for wormholes with a positive total energy density can be found.

From (39), we can obtain the maximum allowed radius of the static metric with the  $\Lambda$  term:

$$r_{\max}^2 = \frac{1 + \sqrt{1 - 4\Lambda q^2}}{2\Lambda} \rightarrow \frac{1}{\Lambda} \text{ as } \Lambda q^2 \rightarrow 0. \tag{43}$$

Beyond this radius, the Universe starts contracting again until a new throat occurs.

Solution (39) describes a static Multiverse with an infinite number of throats. When there is no charge ( $q = 0$ ), this solution becomes a cosmological solution without wormholes, corresponding to a closed isotropic universe (see [31, § 112]).

When there is an excess (or shortage) of the dust part of the energy density or the  $\Lambda$  term relative to (40), the obtained solution for the Multiverse becomes dynamical. Its analytic study is complicated by the need to solve a fourth-order algebraic equation and to calculate a quadrature similar to (37) for the instant of dust layer intersection.

### 8. Conclusion

In this paper, we have generalized and developed the method suggested earlier in papers [20–23, 25] and applied it to new problems in modern cosmology. We found and analyzed analytic solutions of the GR equations describing the dynamics of a traversable wormhole. The results obtained are important for the analysis of the general properties of traversable wormholes. We also obtained a solution describing a spherically symmetric model of the Multiverse. We have not analyzed the properties of geodesics that describe the motion of particles and other matter (energy and information) through a wormhole and its vicinity. These problems, which are mostly important for the analysis of the possible observational appearance of such objects, are considered elsewhere (see, e.g., [11]).

To conclude, we remark on the term ‘Multiverse.’ This term is used in physics and cosmology in two different senses.

First, this term assumes the possibility of the parallel co-existence of many or even an infinite number of different worlds, possibly emerged from a quantum vacuum (in some sense, in different places at different times). In this paper, this term is used exactly in this sense.

Second, this term is sometimes used to denote a sample of different realities in the Everett interpretation of quantum mechanics. The usage of one term for different notions is sometimes confusing. In our opinion, different terms should be used for these concepts. We propose keeping the term ‘Multiverse’ for cosmology, and refer to the set of Everett worlds (following the proposal by M B Mensky) as the ‘Alterverse,’ keeping in mind different classical alternatives of the Everett world.

**Acknowledgments.** We are grateful to the staff of the theoretical astrophysics department of the Lebedev Physical Institute for the discussions. The work is supported by the RFBR grants 07-02-01128a and 08-02-00090a, grants of scientific schools NSh-626.2008.2 and NSh-2469.2008.2, and the program of the Russian Academy of Sciences, The Origin and Evolution of Stars and Galaxies 2008.

## 9. Appendices

### I. Initial conditions with a nonzero velocity

At the WH throat, the expression for  $V^2$  in (26) involves a 0/0 ambiguity. To resolve this ambiguity, we consider the model with a nonzero initial velocity of matter. We introduce the notation

$$1 - \frac{q}{r_i} \equiv \alpha, \quad \frac{r_i}{r} - 1 \equiv \beta, \quad r_{,t}^2|_{(t=0)} \equiv \gamma \tag{44}$$

and assume that  $\Lambda = 0$ . The expressions for the functions  $F_1$  and  $F_2$  with  $\gamma \neq 0$  [see (15) and (12)] are given by

$$F_1 = 1 - s - \frac{q^2}{r_i^2} + \gamma - \frac{\gamma_0 q}{r_i}, \quad F_2 = r_i \left( s + \frac{2q^2}{r_i^2} + \frac{\gamma_0 q}{r_i} \right). \tag{45}$$

With (12) and a nonzero initial velocity, the expression for  $V^2$  in (26) takes the form

$$V^2 = \frac{-s - q^2/r_i^2 + \gamma - \gamma_0 q/r_i + (s + 2q^2/r_i^2 + \gamma_0 q/r_i) r_i/r - q^2/r^2}{1 - s - q^2/r_i^2 + \gamma - \gamma_0 q/r_i}. \tag{46}$$

In the vicinity of the WH throat close to the initial instant, the functions  $\alpha$  and  $\beta$  take small values. From (21), in the linear order in  $\alpha$ , we have

$$s \approx \alpha \varkappa, \quad \varkappa = 8\pi r_i^2 (\varepsilon_i - \varepsilon_d) \approx 8\pi \varepsilon_0 q^2 + 2 < 2.$$

We consider Eqn (46) in the linear order in  $V^2$  with  $\gamma_0 = 0$ . After all transformations, we finally obtain

$$V^2 \approx \frac{\gamma + \gamma_0 \beta}{\gamma - \gamma_0(1 - \alpha) + \alpha(2 - \varkappa)} \rightarrow \frac{\gamma_0(1 + \beta)}{\alpha(\gamma_0 + 2 - \varkappa)} \text{ as } \gamma \rightarrow \gamma_0. \tag{47}$$

It follows that the condition for  $V^2$  to be nonsingular at the throat is that  $\gamma_0 = 0$ . Then

$$V^2 \approx \frac{\gamma}{\gamma + \alpha(2 - \varkappa)}. \tag{48}$$

Expression (48) for  $V^2$  is regular and has no ambiguities if the function  $\gamma$  tends to zero at the throat faster than the function  $\alpha$ . In this case, the rate of change of the radius vanishes at the throat (as must be the case by the definition of the throat). In the linear order in  $\beta$ , the function  $V^2$  is time-independent near the throat.

**II. Study of intersections of adjacent layers**

We consider the model without the  $\Lambda$  term for  $s > 0$ . In this case, the excessive matter collapses, i.e.,  $r \leq r_i$ .

Lemma. *There always exists a nonzero time interval  $[0, t]$  inside which no intersections of adjacent layers occur.*

Equation (30) implies that this is obvious for a nonzero potential. The only point at which this statement must be proved is the point corresponding to  $s = 0$ .

We study the asymptotic regime  $s \rightarrow 0$  in more detail:

$$s \ll \frac{q^2}{r_i^2} \ll 1. \tag{49}$$

From the allowed range of the radius (until the stopping point)  $r \geq q^2/(sr_i + q^2/r_i)$ , we deduce

$$1 - \frac{r}{r_i} \leq \frac{s}{s + q^2/r_i^2} \leq \frac{s}{q^2/r_i^2} \ll 1. \tag{50}$$

Because the last term in (30) contains no  $s$  in the denominator, it can be neglected compared to the previous term. This can be proved in more detail by considering the three exhaustive options for the ratio of the numerator and denominator in the square root expression of the arctan:

1.  $\left(s + \frac{q^2}{r_i^2}\right) \left(1 - \frac{r}{r_i}\right) \ll \left(s + \frac{q^2}{r_i^2}\right) \frac{r}{r_i} - \frac{q^2}{r_i^2}$ .
2.  $\left(s + \frac{q^2}{r_i^2}\right) \left(1 - \frac{r}{r_i}\right) \sim \left(s + \frac{q^2}{r_i^2}\right) \frac{r}{r_i} - \frac{q^2}{r_i^2}$ .
3.  $\left(s + \frac{q^2}{r_i^2}\right) \left(1 - \frac{r}{r_i}\right) \gg \left(s + \frac{q^2}{r_i^2}\right) \frac{r}{r_i} - \frac{q^2}{r_i^2}$ .

Keeping the leading terms, we obtain the asymptotic form of Eqn (30):

$$r' \rightarrow 1 + \frac{(1 - r/r_i)[-q^2 2q^2 2r_i s']}{2sq^4} = 1 - \frac{2r_i s'(1 - r/r_i)}{s}. \tag{51}$$

In a similar way, the first term in the right-hand side of expression (28) can be neglected (compared to the second term), and hence asymptotically with respect to time, we have

$$\begin{aligned} \cos\left(\frac{tq}{r_i^2}\right) &\rightarrow 1 - \frac{2q^2}{sr_i^2} \left(1 - \frac{r}{r_i}\right) \\ &\implies r \rightarrow r_i + \frac{sr_i^3}{2q^2} \left[\cos\left(\frac{tq}{r_i^2}\right) - 1\right]. \end{aligned} \tag{52}$$

Comparing Eqns (51) and (52), we obtain the sought asymptotic form as  $s \rightarrow 0$ :

$$r' \rightarrow 1 - \frac{2r_i^3 s'}{q^2} \sin^2\left(\frac{tq}{2r_i^2}\right). \tag{53}$$

This proves the lemma.

In addition, Eqn (52) implies harmonic dynamics (oscillations) for  $s \rightarrow 0$  and  $2r_i^3 s'/q^2 < 1$ .

**III. Quadrature for  $r'$  in the solution with the  $\Lambda$  term**

We differentiate  $r_{,t}^2$  with respect to  $r_i$ :

$$\frac{dr_{,t}^2}{dr_i} = 2r_{,t} r'_{,t} = 2r_{,t} \frac{dr'}{dr} \implies \frac{dr'}{dr} = \frac{1}{2r_{,t}^2} \frac{dr_{,t}^2}{dr_i}. \tag{54}$$

Substituting Eqn (34) in (54) and differentiating it with respect to  $r_i$ , we obtain the following equation for the function  $y(r) \equiv r'$ :

$$\frac{dy}{dr} + P(r)y = Q(r). \tag{55}$$

Here, the functions  $P(r) \equiv P_1(r) + P_2(r)$  and  $Q(r) \equiv Q_1(r) + Q_2(r)$  are given by (for convenience,  $r_i$  is set to unity below):

$$\begin{aligned} P_1 &= \frac{-1}{2(r-1)}, \\ P_2 &= \frac{1/(2r^2) - (r-1)/r^3 - (a/2)(1 - q^3/r^2)}{(1-r)/r^2 + a(1+r+q^3/r)}, \\ Q_1 &= \frac{-1}{2(r-1)}, \\ Q_2 &= \frac{1/(2r^2) + (r-1)/r^2 + (a/2)(1 - q^3/r)}{(1-r)/r^2 + a(1+r+q^3/r)}. \end{aligned} \tag{56}$$

Equation (55) has a standard solution,

$$\begin{aligned} y(r) &= \exp\left[-\int_{r_1}^r P(x) dx\right] \\ &\times \left\{ \int_{r_2}^r Q(r) \exp\left[+\int_{r_1}^r P(x) dx\right] dr + C_1 \right\}. \end{aligned} \tag{57}$$

The constants  $C_1$ ,  $r_1$ , and  $r_2$  are determined by the initial conditions. The exponential can be rewritten as (the first term with  $P_1$  is integrated)

$$\exp\left[\int_{r_1}^r P(x) dx\right] = \frac{C_2}{\sqrt{r-1}} \exp\left[\int_1^r P_2(x) dx\right]. \tag{58}$$

After that, the part of integral (57) corresponding to  $Q_1$  can also be integrated (by parts), with the result

$$\begin{aligned} y(r) &= \sqrt{r-1} \exp\left[-\int_1^r P_2(x) dx\right] \\ &\times \left\{ \frac{1}{\sqrt{r-1}} \exp\left[+\int_1^r P_2(x) dx\right] \Big|_{r_2}^r \right. \\ &- \int_{r_2}^r \frac{P_2(r)}{\sqrt{r-1}} \exp\left[+\int_1^r P_2(x) dx\right] dr \\ &\left. + \int_{r_2}^r \frac{Q_2(r)}{\sqrt{r-1}} \exp\left[+\int_1^r P_2(x) dx\right] dr + C_1 \right\}. \end{aligned} \tag{59}$$

We can then redefine the lower integration limit and determine the constant  $C_1$  from the initial conditions. The final expression for the function  $y(r) = r'$  is given in (37).

**References**

1. Hawking S W, in *Black Holes and the Structure of the Universe* (Eds C Teitelboim, J Zanelli) (Singapore: World Scientific, 2000) p. 23
2. Lobo F S N *Phys. Rev. D* **71** 084011 (2005)
3. Shinkai H, Hayward S A *Phys. Rev. D* **66** 044005 (2002)
4. Morris M S, Thorne K S *Am. J. Phys.* **56** 395 (1988)
5. Novikov I D *Zh. Eksp. Teor. Fiz.* **95** 769 (1989) [*Sov. Phys. JETP* **68** 439 (1989)]
6. Novikov I D *Phys. Rev. D* **45** 1989 (1992)
7. Hawking S W *Phys. Rev. D* **46** 603 (1992)
8. Visser M *Lorentzian Wormholes: from Einstein to Hawking* (Woodbury, NY: AIP, 1995)
9. Shatskii A A *Astron. Zh.* **81** 579 (2004) [*Astron. Rep.* **48** 525 (2004)]
10. Shatskii A A *Astron. Zh.* **84** 99 (2007) [*Astron. Rep.* **51** 81 (2007)]

11. Kardashev N S, Novikov I D, Shatskii A A *Astron. Zh.* **83** 675 (2006) [*Astron. Rep.* **50** 601 (2006)]
12. Kardashev N S, Novikov I D, Shatskiy A A *Int. J. Mod. Phys. D* **16** 909 (2007)
13. Novikov I D, Kardashev N S, Shatskii A A *Usp. Fiz. Nauk* **177** 1017 (2007) [*Phys. Usp.* **50** 965 (2007)]
14. Cherepashchuk A M *Vestn. Mosk. Gos. Univ., Ser. 3, Fiz. Astron.* (2) 62 (2005) [*Moscow Univ. Phys. Bull.* **60** (2) 74 (2005)]
15. Bronnikov K A, Starobinsky A A *Pis'ma Zh. Eksp. Teor. Fiz.* **85** 3 (2007) [*JETP Lett.* **85** 1 (2007)]; gr-qc/0612032
16. Frolov V P, Novikov I D *Black Hole Physics: Basic Concepts and New Developments* (Fundamental Theories of Physics, Vol. 96) (Dordrecht: Kluwer, 1998)
17. Armendáriz-Picón C *Phys. Rev. D* **65** 104010 (2002); gr-qc/0201027
18. Carr B (Ed.) *Universe or Multiverse?* (Cambridge: Cambridge Univ. Press, 2007)
19. Ellis H G *J. Math. Phys.* **14** 104 (1973)
20. Tolman R C *Proc. Natl. Acad. Sci. USA* **20** 169 (1934)
21. Zel'manov A L *Dokl. Akad. Nauk SSSR* **107** 815 (1956) [*Sov. Phys. Dokl.* **1** 227 (1956)]
22. Novikov I D *Vestnik Mosk. Gos. Univ., Ser. 3, Fiz. Astron.* (5) 90 (1962)
23. Novikov I D *Vestnik Mosk. Gos. Univ., Ser. 3, Fiz. Astron.* (6) 61 (1962)
24. Oppenheimer J R, Snyder H *Phys. Rev.* **56** 455 (1939)
25. Saibal R et al. *Int. J. Mod. Phys. D* **16** 1745 (2007)
26. Shatskii A A, Andreev A Yu *Zh. Eksp. Teor. Fiz.* **116** 353 (1999) [*JETP* **89** 189 (1999)]
27. Shatskii A A *Zh. Eksp. Teor. Fiz.* **131** 851 (2007) [*JETP* **104** 743 (2007)]
28. Fisher I *Zh. Eksp. Teor. Fiz.* **18** 636 (1948)
29. Janis A I, Newman E T, Winicour J *Phys. Rev. Lett.* **20** 878 (1968)
30. Wyman M *Phys. Rev. D* **24** 839 (1981)
31. Landau L D, Lifshitz E M *Teoriya Polya* (The Classical Theory of Fields) (Moscow: Nauka, 1988) [Translated into English (Oxford: Pergamon Press, 1983)]
32. Zel'dovich Ya B, Novikov I D *Relyativistskaya Astrofizika* (Moscow: Nauka, 1967) [Translated into English: *Relativistic Astrophysics* Vol. 1 *Stars and Relativity* (Chicago: Univ. of Chicago Press, 1971)]