

tion of media with noticeable absorption, we used low powers and thin layers of the medium). The radius of the intense beam was several times larger than the dimensions of the particle. The profile of the sounding coloring beam could be chosen arbitrarily.

The display was on a screen located 1.5 m from the layer of the medium. Observation was with the aid of IR binoculars, and motion pictures were taken in the red light of the auxiliary laser.

A growth of the overlap area, by up to 30 times and more, was registered. The strongest effects were observed in Plexiglas, which has a large derivative  $n'_T$  of the refractive index.

We observed<sup>[6]</sup> a new type of self-focusing in the region of perturbations of the medium with  $n'_T > 0$  near the absorbing particle. A smooth rapid transition from the shadow region to the bright point at the center was observed.

These experiments demonstrated the process of aureole refraction from centers giving thermal aureoles with dimensions exceeding the wavelength of the light. The case of aureoles with dimensions that are small compared with the wavelength also admits of a simple description.

The great abundance of natural and artificial media with inhomogeneities (impregnations, sols, dislocations) make the foregoing effects promising in practice. Cases are possible when the transparency and the scattering ability can depend so strongly on the intensity that even a barely noticeable haze or a slightly scattering cloud may turn out to be opaque to light of high intensity. The pulsed character of the processes makes it possible to use them to produce modulators with variable transmission or reflection. These effects can be the cause of power or energy limitation, for example in such working elements as neodymium glass with small platinum particles or other technological impurities.

This effect can be observed also in intense flashes of incoherent light.

<sup>1</sup>G. A. Askar'yan, A. M. Prokhorov, G. F. Chanturiya, and G. P. Shipulo, *Zh. Eksp. Teor. Fiz.* **44**, 2180 (1963) [*Sov. Phys.-JETP* **17**, 1463 (1963)].

<sup>2</sup>G. A. Askar'yan, *ibid.* **45**, 810 (1963) [**18**, 555 (1964)].

<sup>3</sup>O. L. Lebedev and A. A. Chatov, *ibid.* **58**, 848 (1970) [**31**, 455 (1970)].

<sup>4</sup>B. Ya. Kogan and V. L. Churkin, *Opt. Spekr.* **27**, 530 (1969).

<sup>5</sup>Yu. K. Danileiko, A. A. Manenkov, V. S. Nechitaïlo, and V. Ya. Khaimov-Mal'kov, *Zh. Eksp. Teor. Fiz.* **60**, 1245 (1971) [*Sov. Phys.-JETP* **33**, 674 (1971)].

<sup>6</sup>G. A. Askar'yan, V. G. Mikhalevich, and G. P. Shipulo, *ibid.* **60**, 1270 (1971) [**33**, 686 (1971)].

#### B. B. Kadomtsev, Matter in a Superstrong Magnetic Field

In fields of  $10^{12}$ – $10^{24}$  Oe, which according to present notions can exist in neutron stars, there should occur a noticeable change of the physical properties of matter. Namely, a complete realignment of the electron shells

takes place in an atom with atomic number  $Z$  at  $B > Z \times 10^9$  Oe. All the electrons are then at the lower Landau levels, their magnetic moments are oriented along the field, and the electrons move in thin cylindrical shells with symmetry axis passing through the nucleus and directed along the field. In the field interval  $10^9 Z \ll B \ll 10^9 Z^3$  Oe the ground state of the heavy atom can be described within the framework of the modified Thomas-Fermi approximation, which shows that the atom retains its spherical symmetry, and its volume decreases like  $B^{-6/5}$ . In a field  $B \gg 10^9 Z^3$  Oe the atoms are stretched out along the magnetic field, and since they have a large quadrupole moment they can form molecules with high binding energies.

In a sufficiently strong magnetic field, the ionization energy of the atoms and the dissociation energies of the molecules can be so large that the neutral atoms and molecules can exist even at very high temperatures. If the temperatures are not very high, then the atoms and molecules in the superstrong field can become condensed into a solid phase. A preliminary analysis shows that the solid should apparently be a polymer.

<sup>1</sup>B. B. Kadomtsev, *Zh. Eksp. Teor. Fiz.* **58**, 1765 (1970) [*Sov. Phys.-JETP* **31**, 945 (1970)].

<sup>2</sup>B. B. Kadomtsev and V. S. Kudryavtsev, *ZhETF Pis. Red.* **13**, 15, 61 (1971) [*JETP Lett.* **13**, 9, 42 (1971)].

#### I. I. Gurevich, Investigation of the Condensed State of Matter with the Aid of Positive Muons

A new method for studying the properties of method and the kinetics of chemical reactions with the aid of muons has been developed recently at the Atomic Energy Institute, the Nuclear Problems Laboratory of the Joint Institute for Nuclear Research, and the Institute of Theoretical and Experimental Physics. The muon method of investigating the properties of matter is based on the fact that muons stopped in matter are "tagged" particles, the polarization and spin direction of which can be traced by means of the asymmetry of the angular distributions of the  $\mu \rightarrow e$ -decay electrons. It uncovers additional possibilities for studying the kinetics of chemical reactions, local magnetic fields in matter, interactions of muonium atoms with matter, etc.

This paper does not consider interactions of negative muons with matter, which are being intensively studied by V. S. Evseev's group at the Nuclear Problems Laboratory of the Joint Institute for Nuclear Research.

All the experiments considered below on muon interaction with matter were performed with the polarized  $\mu^+$ -meson beam of the same laboratory.

A consistent theory of interaction of positive muons with matter was developed at the Atomic Energy Institute (V. G. Nosov and I. V. Yakovleva, I. G. Ivanter, and V. P. Smilga). According to this theory, when the  $\mu^+$  meson slows down in matter to the velocities of the atomic electrons, it captures an electron and forms a